

SLOWING DOWN AND SPEEDING UP PSR'S PERIODS: A SHAPIRO TELESCOPE TRACING DARK MATTER

D. Fargion, R. Conversano
*Physics Dept. Rome University 1 and INFN,
Piazzale A.Moro 2, 00185 Rome, Italy*

Abstract

Dark matter influence globally galactic Keplerian motions and individually, by microlensing, star's acromatic luminosities. Moreover gravity slows down time as probed by gravitational redshift. Therefore pulsar's (PSR's) periods may record, by slowing and speeding up, any Dark MACHO crossing along the line-of-sight. This phase delay, a Shapiro Phase Delay, is a gravitational integral one and has been probed since 1964 on planets by radar echoes. We discovered in PSRs catalog a few rare PSRs (with negative period derivatives) consistent with speeding up by Shapiro delay due to MACHOs. We propose to rediscover the Shapiro effect on our solar system (for Jupiter, Mars, Sun) to calibrate a new "Shapiro Telescope Array" made by collective monitoring of the sample of PSRs along the ecliptic plane.

1 Introduction

Matter bends space-time. This is the reason for Newton's apple to fall down and for planets and stars to follow Keplerian curves. Gravity is the deformation of space and time. This double reality is why massless photons are bent twice by Einstein Relativity than just one time as predicted by a simplest Newtonian plus Special Relativity model. This factor two is the base for 1919 light deflection test and General Relativity triumph.

The light bending is the root for the microlensing techniques able to reveal unseen dark MACHOs running in our galaxy. But time itself is slowed down by gravity. The famous twin paradox might be played as well as near and far a neutron star surface (neglecting tidal-Riemann forces). The clock beats are slowed down near compact stars (NS, white dwarf, sun..) than in a far flat space- time. Indeed gravitational redshift is an additional cornerstone for General Relativity. The variable position of a wave-source respect to a gravitational field is "recorded" by a phase shift on the wave observed: a Shapiro Phase Delay. Indeed it has been predicted on 1964 and observed by Shapiro

himself with radar echoes from Mercury and Venus, grazing the Sun. Because PSR's period are on average stable ($\langle \dot{P}_{psr} \rangle \leq 10^{-14} \text{ s s}^{-1}$) we proposed [3,4,5] as timing candles to scan all over the galaxy hunting for crossing Dark Matter along the line of sight.

2 The first evidences of Shapiro Phase Delay on PSR's

Microlensing technique seems an unique powerful test because it is inspecting million stars for acromatic luminosity magnifications: how could a few hundred pulsar sample compete with such a huge number? The reason lay on the extreme Shapiro Phase Shift sensitivity versus the microlensing one. While microlensing magnification is proportional to the inverse of the MACHO's impact parameter $\sim 1/b_{min} = 1/u_{min}R_E$ (where u_{min} is adimensional and R_E is the Einstein Radius), Shapiro Phase Delay is made up by two main terms: a first geometrical one, $\Delta t_{geo} = \frac{r_S}{4c} (\sqrt{u^2 + 4} - u)^2$, which is due to deflect and undeflected path and it is rapidly vanishing at large impact values: $\Delta t_{geo}|_{u \gg 1} = \frac{r_S}{cu^2}$. The second and main phase delay is the gravitational redshift (Shapiro Phase Delay) due to a MACHO field $\sim \int \frac{GM}{c^2 r} dr$ all along the wave trajectory; its behaviour is logarithmic on the impact parameter u :

$$\Delta t_{grav} = \frac{r_S}{c} \ln \left(\frac{8D_s}{r_S} \right) - \frac{2r_S}{c} \ln \left(\sqrt{u^2 + 4} + u \right) \quad (1)$$

where $u = b/R_E$ is the adimensional impact parameter of any MACHO crossing along, whose Schwarchild and Einstein radii are:

$$r_S = \frac{GM}{c^2} \quad , \quad R_E = \sqrt{2r_S \frac{D_d D_{ds}}{D_s}} = 4.8 \cdot 10^{13} \sqrt{\left[\frac{M}{M_\odot} \right] \left[\frac{D_s}{4 \text{ Kpc}} \right] \left[\frac{x_d}{1/2} \right] \left[2 - \frac{x_d}{1/2} \right]} \text{ cm} \quad (2)$$

where $x_d = \frac{D_d}{D_s}$, $1 - x_d = \frac{D_{ds}}{D_s}$; D_d is the deflector-observer distance, D_{ds} the source-deflector distance and D_s the source-observer distance.

The consequent characteristic time of the Shapiro delay while the MACHO approaches or goes far away the line of sight is:

$$t_c = \frac{R_E}{v_\perp} u_{min} \simeq 4.55 \frac{\left[\frac{u_{min}}{10^2} \right]}{\left[\frac{\beta_\perp}{10^{-3}} \right]} \sqrt{\left[\frac{M}{M_\odot} \right] \left[\frac{D_s}{4 \text{ Kpc}} \right] \left[\frac{x_d}{1/2} \right] \left[2 - \frac{x_d}{1/2} \right]} \text{ yr} \quad (3)$$

and the consequent PSR's period derivative is:

$$\dot{P} = 1.38 \cdot 10^{-13} \frac{\left[\frac{\beta_{\perp}}{10^{-3}} \right] \sqrt{\frac{M}{M_{\odot}}}}{\left[\frac{u_{min}}{10^2} \right] \sqrt{\left[\frac{D_s}{4 \text{ Kpc}} \right] \left[\frac{x_d}{0.5} \right] \left[2 - \frac{x_d}{0.5} \right]}} s s^{-1} \quad (4)$$

First we must notice that the above period derivative is an order of magnitude larger than the average PSR one ($\dot{P} \sim 10^{-14} s s^{-1}$) and therefore it is well detectable. Secondly because of above sensitivities we could parametrized the adimensional impact parameters not just as $u_{min} \sim 1$ as needed in microlens technique but at $u_{min} \sim 10^2$ scales. This offer a corresponding square geometrical probability amplification (πu^2) to reveal a MACHO, nearly four orders of magnitude larger than in microlensing case. For this reason present Shapiro Delay on 7 hundreds PSRs are comparable with a sample of nearly 7 millions of stars for microlensing. However a positive period derivative ($\dot{P} > 0$) might be well indebted to a large intrinsic positive angular momentum loss. Therefore we first looked for negative PSR period derivative ($\dot{P} < 0$) whose interpretation cannot be indebted to any (rare) corotating accretion mass (single stars). We could find statistically a few events a year [5], for a MACHO density comparable with the observed microlenses. We did observed (up date to 6) and at least one, B1813-26, very isolated PSR, which we do definitively interpret as a Shapiro Phase Delay alive. We also find a group of PSRs in a globular cluster possibly suffering a collective Shapiro Phase Delay due to the own cluster gravitational field: B0021-72C, B0021-72D in 47 Tucanae and B2127+11D, B2127+11A in M15.

3 Shapiro Delay in Dark Planet search

The technique above could be fruitful and certain believed by testing and calibrating the PRS's delay while crossing along the Sun's field, Juppiter or Mars trajectories. For nearby deflectors the gravitational phase delay has a more complicated 3D vectorial formula:

$$\Delta t_{Grav} = \frac{r_S}{c} \ln \left[\frac{D_s - D_d \cos \psi + D_{ds}}{D_d (1 - \cos \psi)} \right] \quad (5)$$

where $D_d = |\vec{D}_d|$, $D_s = |\vec{D}_s|$, $D_{ds} = |\vec{D}_s - \vec{D}_d| = |\vec{D}_{ds}|$ and $\psi = \widehat{\vec{D}_s \vec{D}_d}$. For $\psi \rightarrow 0$ after simple expansion one recovers from equation 5 the previous

Shapiro well known equation 1. For the Sun the effect even few degrees far away its position is huge ($|\dot{P}| \sim 10^{-10} \text{ s s}^{-1}$) and must be observed within $t_c \sim \text{day}$. In order to avoid any additional and confusing refractive index delay (due to solar plasma even at large impact parameters) PRSs might be better observed near Sun at higher radio PSR frequencies [5] ($\nu \geq \text{GHz}$) where plasma refractive index is smaller than the gravitational Shapiro one. Solar Shapiro delay has, naturally, a strong annual modulation. Planets like Jupiter, at 5.2 AU , will produce well detectable period derivative ($|\dot{P}| \sim 10^{-12} \text{ s s}^{-1}$) at $u_{\min} \sim 100$ and even at 10° angular impact parameter distance the effect might be observable ($|\dot{P}| \geq 6 \cdot 10^{-15} \text{ s s}^{-1}$) (see figure 2). In particular to test the technique we chose a sub sample of all known PRSs laying along the ecliptic trajectory ($\pm 6^\circ$). Most PRSs are localized toward Galactic Center and Anticenter and are described in figure 3 [5]. The planetary (Jupiter, Mars..) Shapiro Delay is strongly modulated not only by the Terrestrial yearly trajectory, but also by the combined Earth-Jupiter, Earth-Mars.. mutual distances, which play a key role in defining the Einstein radius and the Shapiro Phase Delay. These combined effects must introduce a very characteristic imprint in their own phase delay as we predicted in figure 3.

4 Conclusions

The Shapiro Phase might be able to discover not only dark matter but once calibrated it may even lead to new mini planets discovers on our ecliptic plane. Their presence might be of relevance (by tidal disturbances during rare encounter) on Earth past (and future) history (and life evolution⁸). Finally recent EGRET data¹⁰ on diffused GeV galactic-Halo might find an answer, among the others⁹, by ($\sim pc$) hydrogen molecular clouds interacting by proton (GeV) cosmic rays¹¹. Microlenses at those large radii, (by Kirchoff theorem) are unefficient. Therefore the Shapiro Phase Delay might be the unique probe to verify such an evanescent dark matter barionic candidature.

1. Alcock C. et al., 1995, SISSA server (astro-ph/9512146)
2. Einstein A., 1936, Sci, 84, 506
3. Fargion D., Conversano R., 1996 , Workshop II “The Dark Side of the Universe”, 252-256.
4. Fargion D., Conversano R., 1997 , MNRAS, 285, 2, 225-231.
5. Fargion D., Conversano R., 1998, astro-ph/9812463
6. Fargion D., 1981, Lett. Nuovo Cim., 31, 49
7. Fargion D., 1983, Lett. Nuovo Cim., 36, 449

8. Fargion D., Dar A., 1988, astro-ph/9802265
9. Fargion D. et al., 1998, astro-ph/9809260
10. Dixon D. D. et al., 1998, astro-ph/9803237 and astro-ph/9803237 v2
11. De Paolis F. et al., 1995, A&A, 295; De Paolis F. et al., 1997, astro-ph/9701214
12. Gebhardt K., 1994, SISSA server, astro-ph/9408086
13. Griest K., 1991, ApJ, 366 ,412
14. King I. R., 1962, AJ, 67, 471
15. King I. R., 1966, AJ, 71, 64
16. Krauss L. M., Small T. A., 1991, ApJ, 378, 22
17. Larchenkova T. I., Doroshenko O.V., 1995, A & A, 297 ,607
18. Madsen J., to appear in MNRAS
19. Manchester R. N., Taylor J. H., 1977, Pulsars. W. H. Freeman and Company, San Francisco
20. Paczyński B., 1986, ApJ, 304 ,1
21. Paczyński B., 1991, ApJ, 374 ,L37
22. Shapiro I. I. , 1964 , Phys. Rev. , 13 , 789
23. Shapiro S. L., Teukolsky S. A., 1983, Black Holes, White Dwarfs, and Neutron Stars, John Wiley & sons, p. 457
24. Schneider P., Ehlers J., Falco E. E., 1992, Gravitational lenses. Springer-Verlag, p. 126
25. Taylor J. H., Manchester R. N., Lyne A. G., 1993, ApJS, 88, 529
26. Taylor J. H., Manchester R. N., Lyne A. G., Camilo F., Princeton Pulsar Group online catalogue.



